

1990




Figure 7.1

SITE: Cowan Creek (Bobbin Head)

SEASON: SUMMER



Habitat




- Seagrass
- Mangroves

Vulnerability of habitats



- Low Vulnerability
- Moderate Vulnerability
- Mod-high Vulnerability
- High Vulnerability

2030



SL=Max




SL=Mean



SL=Min




1990



2050


WT = Mean

AT=Mean




WT = Mean

AT=Mean



WT = Mean

AT=Mean



1990





Figure 7.2

SITE: Cowan Creek (Bobbin Head)


SEASON: WINTER



Habitat






Vulnerability of habitats





2030



SL=Max




SL=Mean



SL=Min




1990



2050


WT = Mean

AT=Mean




WT = Mean

AT=Mean



WT = Mean

AT=Mean



1990




Figure 7.3


SITE: Patonga Entrance

SEASON: SUMMER


N



Habitat






Vulnerability of habitats





2030



SL=Max




SL=Mean



SL=Min




1990



2050


WT = Mean

AT=Mean




WT = Mean

AT=Mean



WT = Mean

AT=Mean



1990






Figure 7.4

SITE: Patonga Entrance


SEASON: WINTER



Habitat






Vulnerability of habitats





2030



SL=Max




SL=Mean



SL=Min




1990



2050


WT = Mean

AT=Mean



WT = Mean

AT=Mean



WT = Mean

AT=Mean

